14-14-14

Item #: 200321-WC

NURSERY POLYON

Homogeneous NPK 4-5 mo @ 70 F.

Average Media Temperatures	60 F	70 F	80 F	90 F
Longevity in Months	5-6	4-5	3-4	2-3

Application Rates

Incorporation

Uniformally incorporate Harrell's Nursery Fertilizer 14-14-14 into potting media at the following rates using trials and grower experience to obtain optimum performance. The addition of dolomitic limestone, additional micronutrients, or other media amendments may be needed

POUNDS PER CUBIC YARD						
Lov 4.5	-	Medium 5.5		High 6.8		

Topdressing - Rates Per Container

Uniformly apply product away from the base of the plant

(gallons)	Low	Medium	High
Trade G (g)	5	6	7
3 (g)	29	36	44
5 (g)	40	48	59
7 (g)	59	72	87
10 (g)	74	90	109
15 (g)	82	100	122
20 (g)	103	126	153



GUARANTEED ANALYSIS				
* Total Nitrogen (N)	14.0000%			
7.3400% Nitrate Nitrogen				
6.6600% Ammoniacal Nitrogen				
** Available Phosphate (P2O5)	14.0000%			
*** Soluble Potash (K2O)	14.0000%			
Calcium (Ca)	2.0000%			

Derived From: Polymer Coated Ammonium Nitrate, Polymer Coated Ammonium Phosphate, Polymer Coated Sulfate of Potash

- 14.3% slow release NITROGEN derived from Polymer Coated Ammonium Nitrate, Polymer Coated Ammonium Phosphate
- ** 14.3% slow release PHOSPHATE derived from Polymer Coated Ammonium Phosphate

*** 14.3% slow release POTASH derived from Polymer Coated Sulfate of Potash

Density - 68 lb./(cu. ft.)

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CONVERSION TABLE		Harrell's SPOON SIZE/RATE		
Rounded Measure	Gram	Spoon Size	Grams	
1 teaspoon (tsp)	7.1		l	
1 tablespoon (tbs.)	22.0	1	10	
1/4 cup	61.4	2	15 20	
1/2 cup	141.5	3 4	20	
[†] The gram weights are	4 5	38		
Actual weight in your r	6	54		
be determined for the fe	7	70		
applied.	0	8	88	
11			-	

KEEP OUT OF REACH OF CHILDREN CAUTION HARMFUL IF SWALLOWED **AVOID INHALATION** OF PRODUCT

APPLICATION PRECAUTIONS

- A trial of this product on a limited scale application methods, and desired rates is recommended. Avoid application to plants under stress.
- Do not incorporate into media prior to steam sterilization.
- For mixes containing soil, reduce application rate by approximately 25%.

When using supplemental liquid feed, reduce the rate of this formulation accordingly.

If mixed media is not used within two weeks, leach thoroughly before using.

To avoid buildup of soluble salts, occasional leaching may be necessary.

MANUFACTURED FOR HARRELL'S LLC. (F352) 720 Kraft Road, Lakeland, FL 33815 - (863) 687-2774 - (800) 282-8007

NET WT Distributed by: Harrell's, LLC. 151 Gene E. Stewart Ct., Sylacauga, AL 35151 - (256) 245-5559 - (800) 282-8007 **50 LBS**

The GREEN Color of the POLYON™ capsule and RLC (Reactive Layer Coating)™ are trademarks of RLC Technologies, L.L.C. POLYON™ and POLYON™ are registered trademarks of Agrium Advanced (22.7 KG) Tochnologios

1410-1309 281410